

## **Colloidal Synthesis of Micron-Scale Organic Crystalline Spherulites**

(No. T4-2334)

## **Principal investigator**

### Dan Oron

Faculty of Chemistry Department of Physics of Complex Systems

### **Principal investigator**

#### Lia Addadi

Faculty of Chemistry Department of Chemical and Structural Biology

## Overview

A controllable method for synthesizing organic, crystalline spherulites from compounds with guanine-based structures. The resulting spherulites are birefringent, having a very high refractive index (RI) in the azimuthal direction and a very low RI in the radial direction, making them excellent broadband scatterers that produce "whiteness." The technology is a way to produce safe alternatives to common pigments such as titanium dioxide, which is a suspected carcinogenic that has been banned as a food additive in the European Union.

## **Applications**

- · Paints and coatings
- Food coloring
- Cosmetics
- Plastics
- Ink and tattoo pigments
- Supplements
- Medicine
- Toothpaste

# Differentiation

- Non-toxic
- Biocompatible
- · Can be applied to a range of materials

# **Development Stage**

The researchers demonstrated the invention with two guanine-based compounds, and a patent was filed. Future technological efforts will focus on biocompatibility tests and exploration of new materials.



Left: Scheme showing the stages of the crystallization process;

Right: SEM image of a broken spherulite produced by mechanical and magnetic stirring

# References

- <u>Nature Photonics 2023</u> [1]
- Advanced Materials 2024 [2]