

Anti-LOX for the Treatment of Duchenne, Fibrosis and Cancer

(No. T4-1913)

Principal investigator

Irit Sagi

Faculty of Biology
Department of Immunology and Regenerative Biology

Overview

LOX is a key driver of fibrosis and tumor progression through its role in extracellular matrix remodeling. We developed a set of LOX inhibitors based on LOX's natural prodomain (LPD) to treat fibrosis, Duchenne Muscular Dystrophy (DMD), and cancer.

Applications


- Treatment of fibrotic diseases and metastatic cancer
- Mutation-independent treatment of DMD

Differentiation

- Selective: targets only extracellular LOX
- Broad: applicable across a range of fibrotic diseases and cancer
- Validated: *in vivo* efficacy in fibrosis and cancer mouse models
- Stable: high-affinity Fc-fusion inhibitor
- Antibody-like: durable and selective design


Development Stage

- Fc- protein inhibitor purified, characterized for stability and high-affinity binding
- Showed significant reduction in fibrosis and improved muscle function in DMD mice
- Reduced lung metastases in a melanoma mouse model



LOX inhibitor

Â



LOX inhibitors (LPD) reduced collagen deposits in the MDX Duchenne mouse model

Patent Status

USA Published: Publication Number: 20240398906 USA Published: Publication Number: 2022-0049236-A1